Amendments to the Claims:

The listing of claims will replace all prior versions, and listings, of claims in the application.

Listing of Claims:

- (original) A method for fostering somatic embryogenic competence of a plant cell
 or tissue comprising contacting said plant cell or tissue with a pro-embryogenic
 arabinosylated protein (AGP) composition comprising a hydrophobic fraction of
 embryogenic AGP and maintaining the cell or tissue in culture to allow the cell or
 tissue to undergo somatic embryogenesis.
- 2. (original) The method of claim 1 wherein the plant cell or tissue is of cotton.
- 3. (currently amended) The method of claim 1 or 2 wherein the plant cell or tissue is selected from the group consisting of Upland cotton, Pima cotton, Egyptian cotton, Sea Island cotton, *G. hirsutum*, *G. barbadens*e, tree cotton, Creole cotton, Levant cotton, Sturt's desert rose cotton, Thurber's cotton, and Hawaii cotton.
- 4. (currently amended) The method of any of claims 1 to 3 claim 1 wherein the plant cell or tissue is of an elite cotton line.
- 5. (currently amended) The method of any of claims 1 to 4 claim 1 wherein the proembryogenic AGP composition comprises embryogenic AGP of a cotton variety.
- 6. (currently amended) The method of any of claims 1 5 claim 1 wherein said AGP composition comprises AGP hydrophobic peak #1 from embryogenic callus from a cotton variety selected from the group consisting of Coker 315, Siokra 1-4, and Sicala 40 at a concentration between about 0.008 and about 0.8 mg/L, and

9

wherein said plant cell or tissue is of a cotton variety that is recalcitrant to

somatic embryogenesis.

7. (original) The method of claim 5 wherein the pro-embryogenic AGP composition

comprises embryogenic AGP selected from the group consisting of de-

glycosylated AGP and de-arabinosylated AGP.

8. (currently amended) The method of claim 5 wherein the pro-embryogenic AGP

is a protein having the amino acid sequence of SEQ ID NO: 25 or SEQ ID NO:

<u> 26</u>.

9. (currently amended) The method of claim 5 wherein the pro-embryogenic AGP

is a thrombin digest of the protein of SEQ ID No: 25 protein having the amino

acid sequence of SEQ ID NO: 26.

10. (currently amended) The method of claim <u>5</u> 8 or 9 wherein the <u>pro-embryogenic</u>

AGP is a thrombin digest of the protein of SEQ ID No: 26 protein is trypsin-

digested.

11. (original) A method for regenerating a plant comprising:

a) harvesting a plant cell or tissue from a first plant;

b) contacting said plant cell or tissue with an AGP composition comprising a

hydrophobic fraction of embryogenic AGP effective for fostering somatic

embryogenic competence; and

c) regenerating a second plant from said plant cell or tissue of step (b).

12. (original) The method of claim 11 comprising, prior to step (b), the step of

transforming said plant cell or tissue whereby a transformed plant is

regenerated.

10

- 13. (currently amended) The method of claim 11 or 12 wherein the plant is cotton.
- 14. (original) The method of claim 13 wherein the cotton plant is a variety selected from the group consisting of Upland cotton, Pima cotton, Egyptian cotton, Sea Island cotton, *G. hirsutum*, *G. barbadens*e, tree cotton, Creole cotton, Levant cotton, Sturt's desert rose cotton, Thurber's cotton, and Hawaii cotton.
- 15. (original) The method of claim 13 wherein the cotton plant is of an elite cotton line.
- 16. (currently amended) The method of any of claims 13 to 15 claim 13 wherein the AGP composition comprises an embryogenic AGP of a cotton variety.
- 17. (original) The method of claim 16 wherein the AGP composition effective for fostering somatic embryogenesis comprises pro-embryogenic AGP selected from the group consisting of de-glycosylated and de-arabinosylated AGP.
- 18. (currently amended) The method of claim 17 wherein the pro-embryogenic AGP is a protein having the amino acid sequence of SEQ ID NO: 25 or SEQ ID NO: 26.
- 19. (currently amended) The method of claim 17 wherein the pro-embryogenic AGP is a thrombin digest of the protein of SEQ ID No: 25 protein having the amino acid sequence of SEQ ID NO: 26.
- 20. (currently amended) The method of claim <u>17</u> 18 or 19 wherein the <u>proembryogenic AGP is a thrombin digest of the protein of SEQ ID No: 26 protein is trypsin-digested.</u>

- 21. (original) A pro-embryogenic AGP composition comprising a hydrophobic fraction of embryogenic AGP of cotton.
- 22. (original) The composition of claim 21 wherein the AGP is de-glycosylated or de-arabinosylated.
- 23. (original) A pro-embryogenic AGP composition comprising a protein comprising a phytocyanin-like domain of an embryogenic AGP.
- 24. (original) A pro-embryogenic AGP composition according to claim 23 comprising a protein having the amino acid sequence of SEQ ID NO: 25.
- 25. (currently amended) The composition of claim <u>23</u> <u>24</u> comprising a <u>thrombin</u> trypsin digest of the protein of SEQ ID No: 25.
- 26. (original) A pro-embryogenic AGP composition according to claim 23 comprising a protein having the amino acid sequence of SEQ ID NO: 26.
- 27. (currently amended) The composition of claim <u>23</u> 26 comprising a <u>thrombin</u> trypsin digest of the protein of SEQ ID No: 25.
- 28. (original) A method for making an AGP composition useful for fostering somatic embryogenic competence comprising:
 - a) providing embryogenic callus; and
 - b) harvesting AGP from said embryogenic callus and fractioning the AGP into hydrophilic and hydrophobic fractions and retaining the hydrophobic fraction.
- 29. (original) A method of making pro-embryogenic AGP by expressing a protein comprising a phytocyanin-like domain of an embryogenic AGP.

- 30. (currently amended) The method of claim 29 wherein the protein has the amino acid sequence of SEQ ID No: 25 or SEQ ID NO:26.
- 31. (canceled)
- 32. (currently amended) The method of claim 29 or 30 comprising the added step of contacting the expressed protein with <u>thrombin</u> trypsin.